

Stability of Leptonic Self-complementarity

Naoyuki Haba¹, Kunio Kaneta^{1,2}, and Ryo Takahashi¹

¹*Department of Physics, Faculty of Science, Hokkaido University, Sapporo 060-0810, Japan and*

²*Department of Physics, Graduate School of Science,
Osaka University, Toyonaka, Osaka 560-0043, Japan*

We investigate a stability of leptonic self-complementarity such that sum of three mixing angles in lepton sector is 90 degrees. Renormalization group equations in a context of minimal supersymmetric standard model for the self-complementarity are analyzed. It is seen that one of Majorana phases plays an important role for the stability of self-complementarity. We find some stable solutions against quantum corrections at a low energy. An effective neutrino mass for neutrino-less double beta decay is also evaluated by the use of neutrino parameters giving rise to the stable solutions.

PACS numbers: 14.60.Pq, 12.60.Jv

Neutrino oscillation experiments established that there are two large mixing angles (θ_{12} and θ_{23}) of Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix in lepton sector. Then a non-vanishing θ_{13} in the PMNS has been reported by recent long baseline and reactor neutrino experiments [1]. These results can be interpreted by three flavor mixing of neutrinos. Regarding with neutrino masses m_i ($i = 1, 2, 3$), the neutrino oscillation experiments determine only two mass squared differences, $\Delta m_{21}^2 \equiv |m_2|^2 - |m_1|^2$ and $|\Delta m_{31}^2| \equiv ||m_3|^2 - |m_1|^2|$. Therefore, two types of neutrino mass hierarchy are allowed, i.e. normal hierarchy (NH) $m_1 < m_2 < m_3$ and inverted hierarchy (IH) $m_3 < m_1 < m_2$. Further, neutrino experiments have not determined whether the neutrinos are Dirac or Majorana particles. Clearly, the nature of neutrinos would be a key to find physics beyond the standard model (SM).

In theoretical side of neutrino physics, various approaches have been discussed in order to investigate hidden flavor structure behind the SM, e.g. introductions of flavor symmetry, mass (matrix) texture analyses, and searches for exotic relations among flavor mixing angles etc.. In this work, we focus on a leptonic self-

complementarity [2] (see also [3] for related discussions) as

$$\theta_{12} + \theta_{23} + \theta_{13} = \frac{\pi}{2} = 90^\circ. \quad (1)$$

We start with effective Yukawa interaction and Weinberg operator at a low energy scale such as electroweak (EW) scale Λ_{EW} in a context of minimal supersymmetric standard model (MSSM), $\mathcal{L}_Y = -y_e \bar{L}_L H_d e_R + \frac{\kappa}{2} (H_u L_L)(H_u L_L) + h.c.$, where L_L are left-handed lepton doublets, e_R are right-handed charged leptons, $H_u(H_d)$ is up(down)-type Higgs, y_e is Yukawa matrix of charged leptons, and $\kappa(H_u L_L)(H_u L_L)$ is the Weinberg operator, which can be effectively induced by integrating out a heavy particle(s). One of examples to obtain this operator is seesaw mechanism. Typical scale of the seesaw mechanism is $\mathcal{O}(10^{14})$ GeV. Therefore, note that the effective coupling κ is having mass dimension -1 and $\kappa^{-1} \sim \mathcal{O}(10^{14})$ GeV. Such a heavy mass scale can realize tiny active neutrino mass scales through the seesaw mechanism. In this work, we utilize an useful parameterization for the PMNS matrix as [4]

$$V_{PMNS} \equiv V_{eL}^\dagger V_\nu D_p = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13} \\ -c_{12}s_{23}s_{13} - s_{12}c_{23}e^{-i\delta} & -s_{12}s_{23}s_{13} + c_{12}c_{23}e^{-i\delta} & s_{23}c_{13} \\ -c_{12}c_{23}s_{13} + s_{12}s_{23}e^{-i\delta} & -s_{12}c_{23}s_{13} - c_{12}s_{23}e^{-i\delta} & c_{23}c_{13} \end{pmatrix} \begin{pmatrix} e^{i\rho} & 0 & 0 \\ 0 & e^{i\sigma} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (2)$$

where $s_{ij} \equiv \sin \theta_{ij}$, $c \equiv \cos \theta_{ij}$ ($i, j = 1, 2, 3$; $i < j$), δ is a Dirac phase, and D_p is a diagonal phase matrix including two Majorana phases, ρ and σ . An neutrino mass matrix M_ν can be diagonalized as $V_\nu^\dagger M_\nu V_\nu^* = M_\nu^{\text{diag}} \equiv \text{Diag}\{m_1, m_2, m_3\}$ with $m_i \equiv \kappa_i v_u^2$ where v_u is vacuum expectation value of up-type Higgs.

Next, we consider renormalization group equations (RGEs) in the MSSM. The RGE of κ is given by $16\pi^2(d\kappa/dt) = \alpha\kappa + [(y_e y_e^\dagger)\kappa + \kappa(y_e y_e^\dagger)^T]$ with $\alpha \equiv 6[-g_1^2/5 - g_2^2 + \text{Tr}(y_u^\dagger y_u)]$ where g_i are gauge coupling

constants, t is an arbitrary renormalization scale as $t \equiv \ln(\mu/\Lambda)$, and Λ is a high energy scale such as the seesaw scale [5, 6]. One can also obtain RGEs of θ_{ij} in a diagonal basis of y_e as $d\theta_{ij}/dt = F_{ij}(\theta_{12}, \theta_{23}, \theta_{13}, \kappa_i, \delta, \rho, \sigma, y_\tau; t)$ where right-hand side (RHS) of this equation is given in [6] (see also [7] for other discussions of mixing angles under the RGEs). Now we turn to the self-complementarity relation (1) and investigate the following equation,

$$\begin{aligned} \frac{d}{dt} \sum_{ij} \theta_{ij} &= \sum_{ij} F_{ij}(\theta_{12}, \theta_{23}, \theta_{13}, \kappa_i, \delta, \rho, \sigma, y_\tau; t) \\ &\equiv F(\theta_{12}, \theta_{23}, \theta_{13}, \kappa_i, \delta, \rho, \sigma, y_\tau; t), \end{aligned} \quad (3)$$

where ij is summed over 12, 23, and 13. The function F is described by 3 mixing angles, 3 effective couplings for the light neutrino masses (or equivalently light neutrino masses m_i), 3 CP-phases, a Yukawa coupling of τ , and renormalization scale. Then once we impose (1) on (3) at an energy scale t_0 , one of mixing angles in F is removed as e.g. $F(\theta_{12}, \theta_{23}, \kappa_i, \delta, \rho, \sigma, y_\tau; t_0)$. We now focus on an equation,

$$\tilde{F}(\theta_{12}, \theta_{23}, \kappa_i, \delta, \rho, \sigma, y_\tau; t_0) = 0. \quad (4)$$

This equation means that once the equation is satisfied at an energy scale t_0 , the self-complementarity is also satisfied at all other energy scales t , i.e. the self-complementarity is stable against quantum corrections, if running effects of parameters except for mixing angles are tiny. In fact, we can find consistent solutions of (4) with experiments for both NH and IH cases. According to the latest experimental data of neutrino oscillation [8]

$$31.3^\circ \lesssim \theta_{12} \lesssim 37.5^\circ, \quad (5)$$

$$38.6^\circ \lesssim \theta_{23} \lesssim 53.1^\circ, \quad (6)$$

$$7.0^\circ (7.3^\circ) \lesssim \theta_{13} \lesssim 10.9^\circ (11.1^\circ), \quad (7)$$

at 3σ level for the NH(IH), the (1) can be satisfied.

Mass spectra of neutrinos at a low energy are defined by $m_1 \equiv \sqrt{m_3^2 - |\Delta m_{31}^2|}$ and $m_2 \equiv \sqrt{m_3^2 - |\Delta m_{31}^2| + \Delta m_{21}^2}$ with best fit values $\Delta m_{21}^2 = 7.62 \times 10^{-5} \text{ eV}^2$ and $\Delta m_{31}^2 = 2.53 \times 10^{-3} \text{ eV}^2$ for the NH, and $m_1 \equiv \sqrt{m_2^2 - \Delta m_{21}^2}$ and $m_3 \equiv \sqrt{m_2^2 - |\Delta m_{31}^2| - \Delta m_{21}^2}$ with $\Delta m_{31}^2 = 2.40 \times 10^{-3} \text{ eV}^2$ for the IH. Therefore, the largest neutrino mass $m_3(m_2)$, of NH(IH) case is a free parameter in our analyses. We analyze in range of $\sqrt{|\Delta m_{31}^2|} \leq m_3 \leq 0.2 \text{ eV}$ ($\sqrt{|\Delta m_{31}^2| + \Delta m_{21}^2} \leq m_2 \leq 0.2 \text{ eV}$). The case of $m_3(m_2) = 0.2 \text{ eV}$ corresponds to a degenerate mass spectrum. In such case, $m_3(m_2)$ is bounded by a cosmological constraint on the sum of neutrino mass as $\sum m_i \lesssim 0.6 \text{ eV}$ [9], and thus as $m_3(m_2) \lesssim \sum m_i/3 \simeq 0.2 \text{ eV}$. Therefore, this must be implied as upper bound on the largest neutrino mass. Here note that since input values of neutrino parameters at a low energy are used, the solutions of (4) are corresponding to ones at $t_0 \simeq \ln(\Lambda_{\text{EW}}/\Lambda)$. The y_τ has been also approximated at a low energy as $y_\tau(\Lambda_{\text{EW}}) = 10^{-2}$ in our analyses.

There are 7 parameters (3 mixing angles, 3 CP-phases, and 1 neutrino mass, m_3 or m_2) and 2 imposed equations ((1) and (4)). Therefore, number of free parameters is 5. Since it is however intricate to deal with all 5 parameters as completely free ones, we numerically analyze at some fixed neutrino masses as examples. According




FIG. 1: Examples of solutions of (4)

to our analyses, a CP-phase is important to give solutions of (4) with (1); we numerically found that there is no solution to satisfy (4) in cases of $(\delta = \rho = \sigma = 0)$, $(\delta \neq 0, \rho = \sigma = 0)$, $(\rho \neq 0, \delta = \sigma = 0)$ for both NH and IH but $(\sigma \neq 0, \delta = \rho = 0)$ can give solutions of (4). In the following, we focus on the case of $(\sigma \neq 0, \delta = \rho = 0)$ in detail. Now we have 2 free parameters (σ and one of mixing angles) in order to look for solution, i.e. once we fix σ and one of mixing angles, all values of our parameters are uniquely determined as we will explain below.

We have scanned over $0 \leq |\sigma| \leq \pi$. Some results of numerical analyses are shown in FIG. 1 as examples. In the figures, vertical and horizontal axes are θ_{23} and θ_{12} , respectively, and shaded region mean that the self-complementarity (1) is correlatively satisfied within 3σ ranges of mixing angles (5)-(7). Lower(upper) slanting and right(left) sides are bounded by maximal(minimal) θ_{13} and θ_{12} at 3σ level, respectively. Note that an allowed range of θ_{23} becomes narrow as $41.6^\circ \lesssim \theta_{23} \lesssim 51.7^\circ$ compared to (6) due to (1). Both two lines in the figures are solutions of (4), which are contours of σ . FIG. 1 (a) is case of $m_3 = \sqrt{|\Delta m_{31}^2|}$ for the NH where $m_1 = 0$, and there are 2 remaining parameters, σ and one of mixing angles. In the case, 2 lines corresponding to $|\sigma| \simeq 126.0^\circ$ and 132.0° are grazing shaded region at upper left described by A and lower right B points, respectively (there are 2 (different sign) solutions of σ in all cases, i.e. solutions are symmetric for reflection respect with π). Therefore, values of all 7 parameters in our analyses are uniquely determined at e.g. point A or B. Further, the points A and B in this case determine maximal and minimal values of parameters for the stability of self-complementarity. Therefore, the self-complementarity is stable in shaded regions of $126.0^\circ \lesssim |\sigma| \lesssim 132.0^\circ$, $31.3^\circ \lesssim \theta_{12} \lesssim 37.5^\circ$, $7.0^\circ \lesssim \theta_{13} \lesssim 10.9^\circ$, and $\text{Min}[\theta_{23}] \leq \theta_{23} \leq \text{Max}[\theta_{23}]$ where $\text{Max}(\text{Min})[\theta_{ij}]$ is maximal(minimal) value of θ_{ij} . $\text{Max}(\text{Min})[\theta_{23}]$ is evaluated by $\text{Max}(\text{Min})[\theta_{23}] = 90^\circ - \text{Min}(\text{Max})[\theta_{12}] - \text{Min}(\text{Max})[\theta_{13}]$ due to (1). Note that value of one of mixing angle (e.g. θ_{23}) is not independently taken because of (1). In the case, we obtain $\text{Max}(\text{Min})[\theta_{23}] \simeq 51.7^\circ (41.6^\circ)$. The results are summarized in TAB. I.

	NH			IH	
m_3	$\sqrt{ \Delta m_{31}^2 }$	$\sqrt{ \Delta m_{31}^2 + \Delta m_{21}^2}$	0.2 eV	0 eV	$\sqrt{m_2^2 - \Delta m_{31}^2 - \Delta m_{21}^2}$
m_2	$\sqrt{\Delta m_{21}^2}$	1.23×10^{-2} eV	0.194 eV	$\sqrt{ \Delta m_{31}^2 + \Delta m_{21}^2}$	5.73×10^{-2} eV
m_1	0 eV	8.73×10^{-3} eV	0.194 eV	$\sqrt{ \Delta m_{31}^2 }$	$\sqrt{m_2^2 - \Delta m_{21}^2}$
Min $[\theta_{12}(\sin^2 \theta_{12})]$	31.3°(0.27)			31.3°(0.27)	
Max $[\theta_{12}(\sin^2 \theta_{12})]$	37.5°(0.37)			37.5°(0.37)	31.5°(0.27)
Min $[\theta_{23}(\sin^2 \theta_{23})]$	41.6°(0.44)			41.4°(0.44)	47.4°(0.54)
Max $[\theta_{23}(\sin^2 \theta_{23})]$	51.7°(0.62)	50.7°(0.60)	51.7°(0.62)	51.4°(0.61)	48.0°(0.55)
Min $[\theta_{13}(\sin^2 \theta_{13})]$	7.0°(0.015)	7.3°(0.016)	7.0°(0.015)	7.3°(0.016)	10.7°(0.034)
Max $[\theta_{13}(\sin^2 \theta_{13})]$	10.9°(0.036)			11.1°(0.037)	
Min $[\sigma]$	126.0°	124.9°	102°	78.8°	87.6°
Max $[\sigma]$	132.0°	π	105°	80.4°	88.8°
Min $[\langle m \rangle_{ee}]$ [meV]	2.36	3.61	73.5	14.9	25.7
Max $[\langle m \rangle_{ee}]$ [meV]	3.09	6.83	97.4	23.1	26.1

TABLE I: Examples of solutions, and minimal and maximal values of neutrino parameters in the corresponding regions

FIG. 1 (b) shows a case of $m_3 = \sqrt{|\Delta m_{31}^2| + \Delta m_{21}^2}$. In the case, contour of $|\sigma| = \pi$, which is the turn of contours of solutions, is across the shaded region and the value of $|\sigma| = \pi$ is a maximal value of $|\sigma|$. Therefore, solutions can be obtained in range of $124.9^\circ \lesssim |\sigma| \leq \pi$. We find that points A-C determine Max(Min) $[\theta_{ij}]$ as $31.3^\circ \lesssim \theta_{12} \lesssim 37.5^\circ$, $7.3^\circ \lesssim \theta_{13} \lesssim 10.9^\circ$, and Max(Min) $[\theta_{23}] \simeq 50.7^\circ(41.6^\circ)$. Derivations of Max $[\theta_{23}]$ and Min $[\theta_{13}]$ are more complicated than the previous case. Max $[\theta_{23}]$ in this case is not determined by Max $[\theta_{23}] = 90^\circ - \text{Min}[\theta_{12}] - \text{Min}[\theta_{13}]$ unlike the previous case because the contour of $|\sigma| = \pi$ does not reach at a upper-left corner of shaded region (this means that the point A does not correspond to Min $[\theta_{13}] \simeq 7.0^\circ$). In this case, the point A where $\theta_{13} \simeq 8.0^\circ$ determines Max $[\theta_{23}]$ as Max $[\theta_{23}] \simeq 90^\circ - \text{Min}[\theta_{12}] - 8.0^\circ \simeq 50.7^\circ$. It is also noticed from the FIG. 1 (b) that the value of Min $[\theta_{13}] \simeq 7.3^\circ$ is not determined at the point A but at C. Therefore, the value of Min $[\theta_{13}] \simeq 7.3^\circ$, which leads to $\theta_{23} \simeq 90^\circ - \text{Max}[\theta_{12}] - 7.3^\circ \simeq 45.2^\circ$, does not determine Max $[\theta_{23}]$ as explained above. Regarding with Max(Min) $[\sigma]$, it corresponds to contours giving Max(Min) $[\theta_{23}]$ in the cases of $m_3 = \sqrt{|\Delta m_{31}^2|}$ and $\sqrt{|\Delta m_{31}^2| + \Delta m_{21}^2}$.

We have also analyzed a case of $m_3 = 0.2$ eV. Results are given in TAB. I. In the case, values of $|\sigma|$ within $102^\circ \lesssim |\sigma| \lesssim 105^\circ$ can give the solutions. Allowed region of mixing angles are the same as ones in the $m_3 = \sqrt{|\Delta m_{31}^2|}$ case, i.e. contours of solutions can reach at the both points A and B like the case of $m_3 = \sqrt{|\Delta m_{31}^2|}$. For the Max(Min) $[\sigma]$, it is determined by Min(Max) $[\theta_{23}]$ in contrast with the above two cases.

We comment on running effects of CP-phases. It has been seen that the one of Majorana phases σ is important for the stability of self-complementarity. One may worry about running effect of σ on low energy solutions for the stability, i.e. whether such effects spoil the solutions at a high energy scale or not. We have approximated the running effects of δ , ρ , and σ from

the seesaw scale on solutions by using leading-log estimation in the RGEs. These running effects $(\Delta\delta, \Delta\rho, \Delta\sigma)$ are $(\Delta\delta, \Delta\rho, \Delta\sigma) \sim (\mathcal{O}(0.001^\circ), \mathcal{O}(0.001^\circ), \mathcal{O}(0.001^\circ))$, $(\mathcal{O}(1^\circ), \mathcal{O}(0.1^\circ), \mathcal{O}(0.1^\circ))$, and $(\mathcal{O}(1^\circ), \mathcal{O}(1^\circ), \mathcal{O}(1^\circ))$ for the cases of $m_3 = \sqrt{|\Delta m_{31}^2|}$, $\sqrt{|\Delta m_{31}^2| + \Delta m_{21}^2}$, and 0.2 eV, respectively. Therefore, most regions of σ are stable against such small running effects up to the seesaw scale because of Max $[\sigma] - \text{Min}[\sigma] > \Delta\sigma$ in all cases. Further, small (but non-vanishing) running effects of δ and ρ do not affect on the stability of self-complementarity. The values in the table do not include such effects of deviations. Then one must remember that our analyses are only for the Majorana neutrino case.

Next, let us consider case of $|\sigma| = \pi$. It is worth emphasizing that all running effects on CP-phases are vanishing in the $|\sigma| = \pi$ case. The solution can be realized in narrow range as $\sqrt{|\Delta m_{31}^2| + \Delta m_{21}^2} \simeq 5.11 \times 10^{-2}$ eV $\lesssim m_3 < 5.15 \times 10^{-2}$ eV. The one of solutions of $|\sigma| = \pi$ is shown in FIG. 1 (b). We find that points A and C constrain mixing angles to $31.3^\circ \lesssim \theta_{12} \lesssim 37.5^\circ$, $7.3^\circ \lesssim \theta_{13} \lesssim 8^\circ$, and Max(Min) $[\theta_{23}] \simeq 50.7^\circ(45.2^\circ)$ in the case of vanishing running effects of CP-phases ($|\sigma| = \pi$).

We have also analyzed IH cases with $m_2 = \sqrt{|\Delta m_{31}^2| + \Delta m_{21}^2}$ and 5.73×10^{-2} eV. The value of $m_2 = 5.73 \times 10^{-2}$ eV is maximal one of m_2 giving solutions. These results are also shown in TAB. I. In the IH case of $m_2 = \sqrt{|\Delta m_{31}^2| + \Delta m_{21}^2}$, there exist solutions in $78.8^\circ \lesssim |\sigma| \lesssim 80.4^\circ$, and mixing angles are constrained to $31.3^\circ \lesssim \theta_{12} \lesssim 37.5^\circ$, $7.3^\circ \lesssim \theta_{13} \lesssim 11.1^\circ$, and Max(Min) $[\theta_{23}] \simeq 51.4^\circ(41.4^\circ)$. For the $m_2 \simeq 5.73 \times 10^{-2}$ eV case, some solutions appear in narrow range as $87.6^\circ \lesssim |\sigma| \lesssim 88.8^\circ$, $31.3^\circ \lesssim \theta_{12} \lesssim 31.5^\circ$, $10.7^\circ \lesssim \theta_{13} \lesssim 11.1^\circ$, and Max(Min) $[\theta_{23}] \simeq 48.0^\circ(47.4^\circ)$.

Finally, we evaluate effective mass term of neutrino-less double beta decay ($0\nu\beta\beta$), $\langle m \rangle_{ee} \equiv |\sum_{i=1}^3 (V_{PMNS})_{ei}^2 m_i|$, in our parameter space. It is written down as

$$\langle m \rangle_{ee} = |m_1 c_{12}^2 c_{13}^2 e^{2i\rho} + m_2 s_{12}^2 c_{13}^2 e^{2i\sigma} + m_3 s_{13}^2|, \quad (8)$$

in our notation. The phenomenon of $0\nu\beta\beta$ can distinguish whether neutrinos are Dirac or Majorana particles. The results at benchmarks given in Tab. I are also presented in the table.

The magnitude of $\langle m \rangle_{ee}$ strongly depends on the scale of m_1 or m_2 rather than mixing angles and CP-phases in the cases. In the NH with $m_3 = \sqrt{|\Delta m_{31}^2|}$ and $\sqrt{|\Delta m_{31}^2| + \Delta m_{21}^2}$, dominant contribution comes from the second term of RHS of (8) because of the small s_{13}^2 and vanishing m_1 . In the cases of NH with $m_3 = 0.2$ eV and IH, the first term of (8) becomes dominant because of mass degeneracy for the NH and inverted mass spectrum for the IH, respectively. We predict 2.36 (14.9) meV $\lesssim \langle m \rangle_{ee} \lesssim 97.4$ (26.1) meV for the NH (IH) within the parameter space to make the self-complementarity stable. The Heidelberg-Moscow experiment [10] for $0\nu\beta\beta$ is giving the most severe bound on $\langle m \rangle_{ee}$, which is $\langle m \rangle_{ee} \lesssim 210$ meV. The CUORE experiment [11] is expected to reach $\langle m \rangle_{ee} = (24 - 93)$ meV in the future. Therefore, a part of our predicting region may be checked in future experiments

We have investigated stability of self-complementarity relation in the PMNS sector against quantum corrections by considering RGEs in the MSSM. We have found solutions stabilizing the self-complementarity both for the NH and IH cases by using low energy data of neutrino oscillation experiments. It has been seen that the one of Majorana phase σ plays an important role to give the solutions. The self-complementarity relation can be satisfied up to an arbitrary high energy scale if neutrino parameters are correlatively within $31.3^\circ \lesssim \theta_{12} \lesssim 37.5^\circ$, $7.0^\circ \lesssim \theta_{13} \lesssim 10.9^\circ$, and $\text{Max}(\text{Min})[\theta_{23}] \simeq 51.7^\circ(41.6^\circ)$ with $126.0^\circ \lesssim |\sigma| \lesssim 132.0^\circ$ and $102^\circ \lesssim \sigma \lesssim 105^\circ$ for the largest neutrino mass $m_3 = \sqrt{|\Delta m_{31}^2|}$ and 0.2

eV of NH with $\delta = \rho = 0$ at a low energy, respectively. For the NH with $m_3 = \sqrt{|\Delta m_{31}^2| + \Delta m_{21}^2}$, we can obtain solutions in regions of $31.3^\circ \lesssim \theta_{12} \lesssim 37.5^\circ$, $7.3^\circ \lesssim \theta_{13} \lesssim 10.9^\circ$, and $\text{Max}(\text{Min})[\theta_{23}] \simeq 50.7^\circ(41.6^\circ)$ with $124.9^\circ \lesssim |\sigma| \lesssim \pi$, $\delta = \rho = 0$. For the IH case with the largest neutrino mass $m_2 = \sqrt{|\Delta m_{31}^2| + \Delta m_{21}^2}$ and $\delta = \rho = 0$ at a low energy, the self-complementarity can be stable in regions of $31.3^\circ \lesssim \theta_{12} \lesssim 37.5^\circ$, $7.3^\circ \lesssim \theta_{13} \lesssim 11.1^\circ$, and $\text{Max}(\text{Min})[\theta_{23}] \simeq 51.4^\circ(41.4^\circ)$. There exist solutions within $\sqrt{|\Delta m_{31}^2|} \leq m_3 \leq 0.2$ eV and $\sqrt{|\Delta m_{31}^2| + \Delta m_{21}^2} \leq m_2 \lesssim 0.0573$ eV for the NH and IH, respectively. In particular, the case of $|\sigma| = \pi$ and $\delta = \rho = 0$, where the RGE running effects on CP-phases are vanishing, predicts $31.3^\circ \lesssim \theta_{12} \lesssim 37.5^\circ$, $7.3^\circ \lesssim \theta_{13} \lesssim 8.0^\circ$, and $\text{Max}(\text{Min})[\theta_{23}] \simeq 50.7^\circ(45.2^\circ)$ for a normal hierarchical neutrino mass spectrum (m_1, m_2, m_3) $\simeq (8.73 \times 10^{-3}, 1.23 \times 10^{-2}, 5.11 \times 10^{-2})$ eV. This kind of solution can be obtained in narrow range as 5.11×10^{-2} eV $\leq m_3 < 5.15 \times 10^{-2}$ eV.

Regarding with the $0\nu\beta\beta$, the effective neutrino mass can be predicted as 2.36 (14.9) meV $\lesssim \langle m \rangle_{ee} \lesssim 97.4$ (26.1) meV for the NH (IH) within the parameter space to make the self-complementarity stable. The above solutions might be checked by combining more precise experimental data of neutrino oscillations, determination of absolute neutrino mass scale, bounds on the sum of neutrino masses, and the discovery of $0\nu\beta\beta$.

Acknowledgment: This work is partially supported by Scientific Grant by Ministry of Education and Science, Nos. 22011005, 24540272, 20244028, and 21244036. The works of K.K. and R.T. are supported by Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists.

-
- [1] K. Abe *et al.* [T2K Collaboration], Phys. Rev. Lett. **107** (2011) 041801; P. Adamson *et al.* [MINOS Collaboration], Phys. Rev. Lett. **107** (2011) 181802; Y. Abe *et al.* [DOUBLE-CHOOZ Collaboration], Phys. Rev. Lett. **108** (2012) 131801; F. P. An *et al.* [DAYA-BAY Collaboration], Phys. Rev. Lett. **108** (2012) 171803; J. K. Ahn *et al.* [RENO collaboration], Phys. Rev. Lett. **108** (2012) 191802.
 - [2] Z. -z. Xing, Phys. Lett. B **696** (2011) 232; Y. -j. Zheng and B. -Q. Ma, Eur. Phys. J. Plus **127** (2012) 7.
 - [3] S. Luo and Z. -z. Xing, arXiv:1203.3118 [hep-ph].
 - [4] H. Fritzsch and Z. -z. Xing, Phys. Lett. B **517** (2001) 363.
 - [5] P. H. Chankowski and Z. Pluciennik, Phys. Lett. B **316** (1993) 312; K. S. Babu *et al.*, Phys. Lett. B **319** (1993) 191.
 - [6] S. Luo *et al.*, Phys. Rev. D **72** (2005) 053014.
 - [7] J. A. Casas *et al.*, Nucl. Phys. B **573** (2000) 652; S. Antusch *et al.*, Nucl. Phys. B **674** (2003) 401; S. Antusch *et al.*, JHEP **0503** (2005) 024; J. -w. Mei, Phys. Rev. D **71** (2005) 073012; N. Haba *et al.*, Prog. Theor. Phys. **103** (2000) 367; J. R. Ellis and S. Lola, Phys. Lett. B **458** (1999) 310. N. Haba *et al.*, Eur. Phys. J. C **10** (1999) 677; Prog. Theor. Phys. **103** (2000) 145; N. Haba and N. Okamura, Eur. Phys. J. C **14** (2000) 347. N. Haba *et al.*, Prog. Theor. Phys. **103** (2000) 807; Eur. Phys. J. C **17** (2000) 513; N. Haba *et al.*, Phys. Lett. B **489** (2000) 184; S. Antusch *et al.*, Nucl. Phys. B **674** (2003) 401; S. Antusch *et al.*, JHEP **0503** (2005) 024; J. -w. Mei, Phys. Rev. D **71** (2005) 073012; J. -w. Mei and Z. -z. Xing, Phys. Lett. B **623** (2005) 227; S. Luo and Z. -z. Xing, arXiv:1203.3118 [hep-ph]; N. Haba and R. Takahashi, arXiv:1206.2793 [hep-ph].
 - [8] D. V. Forero *et al.* arXiv:1205.4018 [hep-ph].
 - [9] E. Komatsu *et al.* [WMAP Collaboration], Astrophys. J. Suppl. **192** (2011) 18.
 - [10] H. V. Klapdor-Kleingrothaus *et al.*, Eur. Phys. J. A **12** (2001) 147.
 - [11] C. Arnaboldi *et al.* [CUORE Collaboration], Nucl. Instrum. Meth. A **518** (2004) 775; M. Sisti [CUORE Collaboration], J. Phys. Conf. Ser. **203** (2010) 012069; F. Bellini *et al.*, Astropart. Phys. **33** (2010) 169.